ELSEVIER

Carbohydrate Research 275 (1995) C13–C16

CARBOHYDRATE RESEARCH

Contents

Regular papers

Separation of cyclodextrins and their derivatives by thin-layer and preparative column chromatography
J. Jindrich, J. Pitha (Baltimore, MD, USA) and B. Lindberg (Stockholm, Sweden) 1

Crystal structures of methyl 6-*O*-acyl-α-D-galactopyranosides
Y. Abe (Ibaraki, Japan), M. Fujiwara, K. Ohbu (Tokyo, Japan) and K. Harata (Ibaraki, Japan). . . . 9

Synthesis of methyl 3,6-dibromo-3,6-dideoxy-β-D-allopyranoside and -glucopyranoside, and their interconversion in the presence of bromide ion
K. Furuhata, N. Aoki, S. Suzuki, N. Arai, H. Ishida (Tokyo, Japan), Y. Saegusa, S. Nakamura (Yokohama, Japan) and M. Sakamoto (Tokyo, Japan) 17

Liquid-crystal properties of octyl 6′-*O*-alkylmelibiosides
L.M. Wingert, G.A. Jeffrey (Pittsburgh, PA, USA), D. Cabaret and M. Wakselman (Versailles, France) 25

Synthesis of 1D-(1,3,5/2,4)-4-acetamido-5-amino-1,2,3- cyclohexanetriol and its incorporation into a pseudo-disaccharide
L.-X. Wang (Shanghai, P.R. China), N. Sakairi (Sapporo, Japan) and H. Kuzuhara (Urawa, Japan) 33

Carbohydrates with relevance to the structure of glycosaminoglycans: The crystal structures of 2-deoxy-2-(sulfoamino)-α-D-glucopyranose sodium salt dihydrate, 2-amino-2-deoxy-α,β-D-glucopyranose 3-(hydrogen sulfate) monohydrate, and 2-amino-2-deoxy-α-D-glucopyranose 6-(hydrogen sulfate) monohydrate
W.H. Ojala, K.E. Albers, W.B. Gleason and C.G. Choo (Minneapolis, MN, USA) 49

Synthesis and structural analysis of disaccharides of 4-*O*-β-D-glucopyranosyl-D-glucosamine and 4-*O*-β-glucopyranosyl-2-deoxy-D-glucose
M.A. Tariq, K. Hayashi, K. Tokuyasu and T. Nagata (Ibaraki, Japan) 67

Crystal structure of anhydrous heptakis-(2,6-di-*O*-methyl)cyclomaltoheptaose (dimethyl-β-cyclodextrin)
T. Steiner and W. Saenger (Berlin, Germany). 73

Stereoselective synthesis of L-vancosamine methyl β-glycoside by addition of an organocerium reagent to *O*-benzyloxime ethers
R. Greven, P. Jütten and H.-D. Scharf (Aachen, Germany). 83

Synthesis and reactions of 1-(5-azido-5-deoxy-3-*O*-*p*-toluenesulfonyl-β-D-xylofuranosyl) derivatives of 5-alkyl- and 5-halo-pyrimidines
N.A. Al-Masoudi (Basrah, Iraq) and W. Pfleiderer (Konstanz, Germany) 95

β-Elimination of protected aldono-1,4-lactones as a general approach to the synthesis of 2-keto-3-deoxyaldonic acids containing four to six carbon atoms
G. Limberg and J. Thiem (Hamburg, Germany) 107

Synthesis and crystal structure of methyl 4,6-dideoxy-4-(3-deoxy-L-*glycero*-tetronamido)- 2-*O*-methyl-α-D-mannopyranoside, the methyl α-glycoside of the terminal unit, and presumed antigenic determinant, of the O-specific polysaccharide of *Vibrio cholerae* O:1, serotype Ogawa
P. Lei, Y. Ogawa (Bethesda, MD, USA), J.L. Flippen-Anderson (Washington, DC, USA) and P. Kováč (Bethesda, MD, USA) 117

Chain conformation in concentrated pectic gels: evidence from ^{13}C NMR
M.C. Jarvis (Glasgow, UK) and D.C. Apperley (Durham, UK) 131

Structure of the capsular polysaccharide from *Alteromonas sp.* CMM 155
V.A. Zubkov, E.L. Nazarenko, R.P. Gorshkova, E.P. Ivanova (Vladivostok, Russian Federation), A.S. Shashkov, Y.A. Knirel, N.A. Paramonov (Moscow, Russian Federation) and Y.S. Ovodov (Vladivostok, Russian Federation) 147

Structure of the repeating units in the rhamnogalacturonic backbone of apple, beet and citrus pectins
C.M.G.C. Renard, M.-J. Crépeau and J.-F. Thibault (Nantes, France) 155

Lignin–ferulate cross-links in grasses: active incorporation of ferulate polysaccharide esters into ryegrass lignins
J. Ralph, J.H. Grabber and R.D. Hatfield (Madison, WI, USA) 167

Notes

Synthesis of 1-thioglycosides
P. Li, L. Sun, D.W. Landry and K. Zhao (New York, NY, USA) 179

Complete assignments of ^{13}C NMR resonances to all the carbon atoms of the trimannosido-di-*N*-acetylchitobiosyl structure in a pentaantennary decasaccharide glycopeptide
T. Taguchi, K. Kitajima, T. Niimi, Y. Muto, S. Yokoyama, S. Inoue and Y. Inoue (Tokyo, Japan) . 185

Isolation and characterization of a new β-galactose-specific lectin from the sea worm *Chaetopterus variopedatus*
L.V. Mikheyskaya, E.V. Evtushenko, R.G. Ovodova, N.I. Belogortseva and Y.S. Ovodov (Vladivostok, Russian Federation) 193

The structure of the O-specific polysaccharide chain of *Proteus penneri* strain 42 lipopolysaccharide
Y.A. Knirel, A.S. Shashkov, E.V. Vinogradov (Moscow, Russian Federation), A. Swierzko and Z. Sidorczyk (Lodz, Poland) 201

Purification and characterization of a β-mannanase from the digestive tract of the edible snail *Helix lucorum* L.
V. Flari (Thessaloniki, Greece), M. Matoub and C. Rouland (Créteil, France). 207

Differential activities of a lipase and a protease toward straight- and branched-chain acyl donors in transesterification to carbohydrates in an organic medium
T. Watanabe, R. Matsue, Y. Honda and M. Kuwahara (Kyoto, Japan). 215

Regular papers

Synthesis of pentasaccharide analogues of the *N*-glycan substrates of *N*-acetylglucosaminyltransferases III, IV and V using tetrasaccharide precursors and recombinant β-(1 → 2)-*N*-acetylglucosaminyltransferase II
F. Reck (Toronto, Canada), E. Meinjohanns (Hamburg, Germany), J. Tan, A.A. Grey (Toronto, Canada), H. Paulsen (Hamburg, Germany) and H. Schachter (Toronto, Canada) 221

Synthetic mucin fragments: synthesis of *O*-sulfo and *O*-methyl derivatives of allyl *O*-(β-D-galactopyranosyl)-(1 → 3)-2-acetamido-2-deoxy-α-D-galactopyranoside as potential compounds for sulfotransferases
R.K. Jain, C.F. Piskorz and K.L. Matta (Buffalo, NY, USA) 231

Spin-labelled glycolipid analogues: D-glucose series
J.M.J. Tronchet, M. Zsély and M. Geoffroy (Geneva, Switzerland) 245

Synthesis, conformation and glycosylation reaction of substituted 2,6-dioxabicyclo[3.1.1]heptanes: 1,3-anhydro-2,4-di-*O*-benzyl-α-L-arabinopyranose
Y. Du and F. Kong (Beijing, P.R. China) 259

The N-7 regioisomer of 2-chloro-2′-deoxyadenosine: synthesis, crystal structure, conformation, and stability
V.L. Worthington, W. Fraser and C.H. Schwalbe (Birmingham, UK) 275

Structural analysis of the O-antigen of the lipopolysaccharide of *Rhizobium tropici* CIAT899
A.M. Gil-Serrano, I. González-Jiménez, P. Tejero Mateo (Seville, Spain), M. Bernabé, J. Jiménez-Barbero (Madrid, Spain), M. Megías and M.J. Romero-Vázquez (Seville, Spain) 285

Characterization of a monoclonal antibody that recognizes an arabinosylated (1 → 6)-β-D-galactan epitope in plant complex carbohydrates
W. Steffan (Athens, GA, USA), P. Kovář (Bethesda, MD, USA), P. Albersheim, A.G. Darvill and M.G. Hahn (Athens, GA, USA) 295

Analysis of cationic starches: determination of the substitution pattern of *O*-(2-hydroxy-3-trimethylammonium)propyl ethers
O. Wilke and P. Mischnick (Hamburg, Germany) 309

Structure of an exocellular polysaccharide of *Lactobacillus helveticus* TN-4, a spontaneous mutant strain of *Lactobacillus helveticus* TY1-2
Y. Yamamoto (Gifu, Japan), T. Nunome (Hyougo, Japan), R. Yamauchi, K. Kato (Gifu, Japan) and Y. Sone (Osaka, Japan) 319

Structural elucidation of the capsular polysaccharide of *Bacteroides fragilis* strain 23745M1
V. Pavliak, D. Uhrín, J.-R. Brisson (Ottawa, Canada), A.O. Tzianabos, D.L. Kasper (Boston, MA, USA) and H.J. Jennings (Ottawa, Canada) 333

Different populations of pectic hairy regions occur in apple cell walls
H.A. Schols, E. Vierhuis, E.J. Bakx and A.G.J. Voragen (Wageningen, The Netherlands) 343

Structural analysis of starch from normal and *shx* (shrunken endosperm) barley (*Hordeum vulgare* L.)
A.H. Schulman (Helsinki, Finland), S. Tomooka (Kagoshima, Japan), A. Suzuki (Sakai, Japan), P. Myllärinen (Helsinki, Finland) and S. Hizukuri (Kagoshima, Japan) 361

The structure of the acidic exopolysaccharide produced by *Pseudomonas* "gingeri" strain Pf9
P. Cescutti, S.F. Osman, W.F. Fett (Philadelphia, PA, USA) and D. Weisleder (Peoria, IL, USA) . . 371

Notes

Gradient-enhanced homonuclear 2D NMR techniques applied to oligosaccharides containing mannohexoses provide improved correlations for protons coupled by small ^3J
A. Otter, O. Hindsgaul and D.R. Bundle (Edmonton, Canada) 381

Conformational analysis of sucrose octasulfate by high resolution nuclear magnetic resonance spectroscopy
U.R. Desai, I.R. Vlahov, A. Pervin and R.J. Linhardt (Iowa City, IA, USA) 391

Synthese von modifizierten Tetrasacchariden als analoge Akzeptor–Inhibitoren der *N*-Acetylglucosaminyltransferase II
H. Paulsen, M. Springer (Hamburg, Deutschland), F. Reck, I. Brockhausen and H. Schachter (Toronto, Canada) 403

Synthesis and unusual glycosidic coupling reaction of substituted 2,7-dioxabicyclo[4.1.0]heptanes: 1,2-anhydro-3,4-di-*O*-benzyl-α-D-fucopyranose
Y. Du and F. Kong (Beijing, P.R. China) 413

Regioselective chlorination study of unprotected pentitols: syntheses of some 1,5-dichloro-1,5-dideoxypentitol derivatives
M. Benazza, M. Massoui (Kénitra, Morocco), R. Uzan and G. Demailly (Amiens, France). 421

Inactivation of two D-glucosyltransferases from Serotype c *Streptococcus mutans* 103220T by 1,2-epoxy-3-(α-D-glucopyranosyl)propane
S. Bombard, M. Maillet and M.-L. Capmau (Paris, France) 433

Author index C3

Subject index C5

Contents C13